

DGL SYNERGY™

FAST ACTING. CHEWABLE DEGLYCYRRHIZINATED LICORICE FOR SUPPORT OF THE GASTRIC MUCOSA 90 TABLETS | NPN80057886 | DGL090-CN

DGL Synergy™ is a pleasant tasting, chewable formula of deglycyrrhizinated licorice (DGL), to help with dyspepsia and symptoms associated with gastroesophageal reflux disease, or GERD. It contains a standardized high-guality form of DGL, the amino acid glycine, and calcium in the form of calcium glycerophosphate.

DGL (Glycyrrhiza glabra) root extract - Well-recognized for its anti-ulcer and anti-inflammatory properties, which exhibit effects upon the gastric mucosa. Licorice is a mucilaginous herb that coats and soothes inflamed gastric and intestinal tissue. With its very low glycyrrhizin content, DGL works efficaciously without some of the potential side effects of prolonged high-dose full-spectrum licorice, including hypertension, hypokalemia (low blood potassium levels) and fluid retention. DGL is also potent for attenuating H. pylori, an organism which, when overgrown, is associated with gastritis, ulcers and tumorigenesis.

- May help reduce aspirin-induced damage to gastric tissue^{1,2,3}
- Promotes secretion of protective gastric mucus and may prolong the lifespan of mucus-producing cells⁴
- Has shown effectiveness for aphthous ulcers (mouth sores, canker sores) and inflamed oral tissue⁵

Glycine - Conditionally essential amino acid required for synthesis of bile acids; may help in proper digestion of dietary fats9

Calcium Glycerophosphate - Included in this product for its acid-reducing properties in the gut¹⁰

BENEFITS

Effective for both acute and chronic symptoms of GERD, including:

Dyspepsia

Heartburn

Acid reflux

HIGHLIGHTS OF DGL SYNERGY™

- Fast-acting, chewable tablet Great licorice flavour
 - Made with GMO-free ingredients
- Fructose-free and sugar-free; sweetened with the polyols xylitol and mannitol, and the plant extract luo han guo

Medicinal Ingredients (per tablet):

DGL (Deglycyrrhizinated licorice, Glycyrrhiza glabra-Root) (1.004:1)	250 mg
Glycine (Aminoacetic acid)	100 mg
Calcium (Calcium glycerophosphate)	

Non-Medicinal Ingredients: Xylitol, d-mannitol, cellulose, magnesium stearate (vegetable source), Siraitia grosvenorii fruit extract. Recommended Dose: Adults: Take 1 tablet 3 times per day. Chew between meals or 20 minutes before meals. Take a few hours before or after taking other medications.

REFERENCES

- 1. Effect of deglycyrrhizinated liquorice on gastric mucosal damage by aspirin. Rees WD, Rhodes J, Wright JE, Stamford LF, Bennett A. Scand J Gastroenterol. 1979;14(5):605-7.
- 2. The protective effect of liquorice components and their derivatives against gastric ulcer induced by aspirin in rats. Dehpour AR. Zolfaghari ME. Samadian T. Vahedi Y. J Pharm Pharmacol. 1994 Feb: 46(2):148-9.
- 3. Deglycyrrhizinised liquorice: a report of its pharmacological properties and ther—apeutic efficacy in peptic ulcer. Brogden RN, Speight TM, Avery GS. Drugs. 1974;8(5):330-9.
- 4. Licorice: A possible anti-inflammatory and anti-ulcer drug. Aly AM, Al-Alousi L, Salem HA. AAPS PharmSciTech. Mar 2005; 6(1): E74-E82.
- 5. Deglycyrrhizinated liquorice in aphthous ulcers. Das SK, Das V, Gulati AK, Singh VP. J Assoc Physicians India. 1989 Oct;77(10):647.
- 6. Effect of licorice versus bismuth on eradication of Helicobacter pylori in patients with peptic ulcer disease. Momeni A, Rahimian G1, Kiasi A1, Amiri M2, Kheiri S3. Pharmacognosy Res. 2014 Oct;6(4):341-4.
- 7. Special licorice extracts containing lowered glycyrrhizin and enhanced licochal—cone a prevented Helicobacter pylori-initiated, salt diet-promoted gastric tumori—genesis. Park JM, Park SH, Hong KS, Han YM, Jang SH, Kim EH, Hahm KB. Helicobacter. 2014 Jun;19(3):221-36.
- 8. Anti-Helicobacter pylori flavonoids from licorice extract. Fukai T, Marumo A, Kaitou K, Kanda T, Terada S, Nomura T. Life Sci. 2002 Aug 9;71(12):1449-63.
- 9. Glycine metabolism in animals and humans: implications for nutrition and health. Wang W, Wu Z, Dai Z, Yang Y, Wang J, Wu G. Amino Acids. 2013 Sep;45(3):463-77.
- 10. The efficacy of calcium glycerophosphate in the prevention of food-related flares in interstitial cystitis. Bologna RA, Gomelsky A, Lukban JC, Tu LM, Holzberg AS, Whitmore KE. Urology. 2001 Jun;57(6 Suppl 1):119-20.